

METHOD FOR PRODUCING XYLYLENEDIAMINE (XDA)

Publication number: WO2005028417

Publication date: 2005-03-31

Inventor: HUGO RANDOLF (DE); JOURDAN SABINE (DE); WENZ KIRSTEN (DE); PREISS THOMAS (DE); WECK ALEXANDER (DE)

Applicant: BASF AG (DE); HUGO RANDOLF (DE); JOURDAN SABINE (DE); WENZ KIRSTEN (DE); PREISS THOMAS (DE); WECK ALEXANDER (DE)

Classification:

- international: **C07C209/48; C07C211/27; C07C209/00; C07C211/00;**
(IPC1-7): C07C209/48; C07C211/27

- European: C07C209/48

Application number: WO2004EP09568 20040827

Priority number(s): DE20031041613 20030910

Also published as:

EP1663942 (A1)
EP1663942 (A0)
DE10341613 (A1)
CN1849295 (A)

Cited documents:

EP1193244
US2970170
EP1279661
GB1164354
EP1209146

Report a data error here

Abstract of WO2005028417

The invention relates to a method for producing xylylenediamine comprising the steps: ammoxidation of xylol to form phthalodinitrile, the vaporous product of the ammoxidation step being brought into direct contact with a liquid, organic solvent (quench stage) and hydrogenation of the phthalodinitrile in the quench solution or suspension that has been obtained, whereby N-methyl-2-pyrrolidone (NMP) constitutes the organic solvent.

